

Physics
Standard level
Paper 2

Friday 17 May 2019 (afternoon)

Candidate session number

1 hour 15 minutes

--	--	--	--	--	--	--	--	--	--

Instructions to candidates

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Answer all questions.
- Answers must be written within the answer boxes provided.
- A calculator is required for this paper.
- A clean copy of the **physics data booklet** is required for this paper.
- The maximum mark for this examination paper is **[50 marks]**.



Answer **all** questions. Answers must be written within the answer boxes provided.

1. A student strikes a tennis ball that is initially at rest so that it leaves the racquet at a speed of 64 m s^{-1} . The ball has a mass of 0.058 kg and the contact between the ball and the racquet lasts for 25 ms .

(a) Calculate the

- (i) average force exerted by the racquet on the ball.

[2]

.....

.....

.....

.....

- (ii) average power delivered to the ball during the impact.

[2]

.....


.....

.....

.....

- (b) The student strikes the tennis ball at point P. The tennis ball is initially directed at an angle of 7.00° to the horizontal.

diagram not to scale



The following data are available.

Height of P	$= 2.80 \text{ m}$
Distance of student from net	$= 11.9 \text{ m}$
Height of net	$= 0.910 \text{ m}$
Initial speed of tennis ball	$= 64 \text{ m s}^{-1}$

(This question continues on the following page)



(Question 1 continued)

- (i) Calculate the time it takes the tennis ball to reach the net. [2]

.....

.....

.....

.....

- (ii) Show that the tennis ball passes over the net. [3]

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) Determine the speed of the tennis ball as it strikes the ground. [2]

.....

.....

.....

.....

.....


.....

(This question continues on the following page)



(Question 1 continued)

- (c) The student models the bounce of the tennis ball to predict the angle θ at which the ball leaves a surface of clay and a surface of grass.



The model assumes

- during contact with the surface the ball slides.
- the sliding time is the same for both surfaces.
- the sliding frictional force is greater for clay than grass.
- the normal reaction force is the same for both surfaces.

Predict for the student's model, without calculation, whether θ is greater for a clay surface or for a grass surface.

[3]

.....

.....

.....

.....

.....

.....



2. A container of volume $3.2 \times 10^{-6} \text{ m}^3$ is filled with helium gas at a pressure of $5.1 \times 10^5 \text{ Pa}$ and temperature 320 K . Assume that this sample of helium gas behaves as an ideal gas.

- (a) The molar mass of helium is 4.0 g mol^{-1} . Show that the mass of a helium atom is $6.6 \times 10^{-27} \text{ kg}$.

[1]

.....

.....

- (b) Estimate the average speed of the helium atoms in the container.

[2]

.....

.....

.....

.....

- (c) Show that the number of helium atoms in the container is about 4×10^{20} .

[2]

.....

.....

.....

.....

- (d) A helium atom has a volume of $4.9 \times 10^{-31} \text{ m}^3$.

- (i) Estimate the ratio $\frac{\text{total volume of helium atoms}}{\text{volume of helium gas}}$.

[1]

.....

.....

- (ii) Explain, using your answer to (d)(i) and with reference to the kinetic model, why this sample of helium can be assumed to be an ideal gas.

[2]

.....

.....

.....

.....



3. The diagram shows the direction of a sound wave travelling in a metal sheet.




diagram not to scale

- (a) Particle P in the metal sheet performs simple harmonic oscillations. When the displacement of P is $3.2\ \mu\text{m}$ the magnitude of its acceleration is $7.9\ \text{ms}^{-2}$. Calculate the magnitude of the acceleration of P when its displacement is $2.3\ \mu\text{m}$. [2]

.....

.....

.....

.....

- (b) The wave is incident at point Q on the metal–air boundary. The wave makes an angle of 54° with the normal at Q. The speed of sound in the metal is $6010\ \text{ms}^{-1}$ and the speed of sound in air is $340\ \text{ms}^{-1}$. Calculate the angle between the normal at Q and the direction of the wave in air. [2]

.....

.....

.....

.....

(This question continues on the following page)



(Question 3 continued)

(c) The frequency of the sound wave in the metal is 250 Hz.

(i) State the frequency of the wave in air.

[1]

.....

.....


(ii) Determine the wavelength of the wave in air.

[1]

.....

.....

(d) The sound wave in air in (c) enters a pipe that is open at both ends. The diagram shows the displacement, at a particular time T , of the standing wave that is set up in the pipe.




On the diagram, at time T , label with the letter C a point in the pipe that is at the centre of a compression.

[1]



4. Three identical light bulbs, X, Y and Z, each of resistance $4.0\ \Omega$ are connected to a cell of emf 12 V. The cell has negligible internal resistance.



- (a) The switch S is initially open. Calculate the total power dissipated in the circuit.

[2]

.....

.....

.....

.....

- (b) The switch is now closed.

- (i) State, without calculation, why the current in the cell will increase.

[1]

.....

.....

.....

- (ii) Deduce the ratio $\frac{\text{power dissipated in Y with S open}}{\text{power dissipated in Y with S closed}}$

[2]

.....


.....

.....

.....



5. A proton moves along a circular path in a region of a uniform magnetic field. The magnetic field is directed into the plane of the page.



- (a) Label with arrows on the diagram the
- (i) magnetic force F on the proton. [1]
 - (ii) velocity vector v of the proton. [1]
- (b) The speed of the proton is $2.16 \times 10^6 \text{ m s}^{-1}$ and the magnetic field strength is 0.042 T . For this proton, determine, in m, the radius of the circular path. Give your answer to an appropriate number of significant figures. [3]

.....

.....

.....

.....

.....

.....





(This question continues on the following page)

.....
.....
.....
.....

(ii) Suggest why, for the fusion reaction above to take place, the temperature of deuterium must be very high. [2]

.....
.....
.....
.....
.....
.....


(i) Determine, in MeV, the energy released. [2]

$${}^1_2\text{H} = 1.12\text{MeV}$$
$${}^1_3\text{H} = 2.78\text{MeV}$$

(b) The following data are available for binding energies per nucleon.

.....
.....


(a) Identify particle X. [1]



6. Deuterium, ${}^1_2\text{H}$, undergoes fusion according to the following reaction.

(Question 6 continued)

- (c) Particle Y is produced in the collision of a proton with a K^- in the following reaction.



The quark content of some of the particles involved are

$$K^- = \bar{u}s \quad K^0 = d\bar{s}$$

Identify, for particle Y, the

- (i) charge.

[1]

.....
.....

- (ii) strangeness.

[1]

.....
.....



7. The average temperature of ocean surface water is 289 K. Oceans behave as black bodies.

(a) Show that the intensity radiated by the oceans is about 400 W m^{-2} . [1]

.....

.....

(b) Explain why some of this radiation is returned to the oceans from the atmosphere. [3]

.....

.....

.....

.....

.....

.....

