



Physics Standard level Paper 3

2 May 2023

Zone A afternoon | Zone B morning | Zone C morning

Candidate session number

1 hour

--	--	--	--	--	--	--	--	--	--

Instructions to candidates

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Answers must be written within the answer boxes provided.
- A calculator is required for this paper.
- A clean copy of the **physics data booklet** is required for this paper.
- The maximum mark for this examination paper is **[35 marks]**.

Section A	Questions
Answer all questions.	1 – 2

Section B	Questions
Answer all of the questions from one of the options.	
Option A — Relativity	3 – 5
Option B — Engineering physics	6 – 7
Option C — Imaging	8 – 10
Option D — Astrophysics	11 – 13




Section A

Answer **all** questions. Answers must be written within the answer boxes provided.

1. A student investigates the relationship between the pressure in a ball and the maximum force that the ball produces when it rebounds.

A pressure gauge measures a difference Δp between the atmospheric pressure and the pressure in the ball. A force sensor measures the maximum force F_{\max} exerted on it by the ball during the rebound.



- (a) State **one** variable that needs to be controlled during the investigation.

[1]

(This question continues on the following page)



(Question 1 continued)

The student collects the following data.

Gauge pressure Δp / kPa	Maximum force F_{\max} / N
10	108
20	133
30	158
40	170
50	188
60	192
70	206
80	220

The student initially hypothesizes that Δp is proportional to F_{\max} .

- (b) Deduce, using **two** suitable data points from the table, that the student's initial hypothesis is not supported.

[3]

.....

.....

.....

.....

.....

.....


(This question continues on the following page)



(Question 1 continued)

The student now proposes that $F_{\max}^3 = k\Delta p$.

The student plots a graph of the variation of Δp with F_{\max}^3 .



(c) (i) State the unit for k .

[1]

.....

(ii) Plot on the graph the position of the missing point for the Δp value of 40 kPa.

[1]

(This question continues on the following page)



(Question 1 continued)

The percentage uncertainty in F_{max} is $\pm 5\%$. The error bars for F_{max}^3 at $\Delta p = 10 \text{ kPa}$ and $\Delta p = 80 \text{ kPa}$ are shown.

- (d) (i) Calculate the absolute uncertainty in F_{max}^3 for $\Delta p = 30 \text{ kPa}$. State an appropriate number of significant figures for your answer.

[3]

.....

.....

.....

.....

.....

.....

- (ii) Plot the absolute uncertainty determined in part (d)(i) as an error bar on the graph. [1]

- (iii) Explain why the new hypothesis is supported. [1]

.....

.....

.....

.....



2. A student conducts an experiment to determine the specific heat capacity of a metal cube. The cube is heated in a beaker of boiling water to a temperature of 100°C and then quickly transferred into an insulated vessel of negligible thermal capacity. The vessel contains water at 20°C and of known specific heat capacity.

(a) State one other measurement that the student will need to make.

[1]

.....

(b) Suggest one modification that the student can make to reduce the fractional uncertainty for the change in temperature of the metal cube.

[1]

.....

(c) Some water from the beaker is accidentally transferred with the cube.

Discuss how this will affect the value of the calculated specific heat capacity of the cube. [2]

.....




Section B

Answer **all** of the questions from **one** of the options. Answers must be written within the answer boxes provided.

Option A — Relativity

3. A wire carries an electric current. An external electron e moves with the drift velocity v of the electrons in the wire. Observer O is at rest relative to the wire.



- (a) State what is meant by a frame of reference.

[1]

- (b) State and explain the nature of the electromagnetic force acting on electron e in the frame of reference of

- (i) observer O .

[2]

- (ii) electron e .

[2]

(Option A continues on the following page)




32EP07

Turn over

(Option A continued)

4. Star A and star B are separated by a fixed distance of 4.8 light years as measured in the reference frame in which they are stationary. An observer P at rest in a space station moves to the right with speed $0.78c$ relative to the stars. A shuttle S travels from star A to star B at a speed of $0.30c$ relative to the stars.



- (a) State the value of the maximum distance between the stars that can be measured in any reference frame. [1]

.....

.....

- (b) Write down the speed of shuttle S relative to observer P using Galilean relativity. [1]

.....

.....

- (c) Calculate the distance between star A and star B relative to observer P. [2]

.....

.....

.....

.....

(Option A continues on the following page)



(Option A, question 4 continued)

- (d) Show that the speed of shuttle S relative to observer P is approximately $0.6c$. [2]

.....

.....

.....

.....

- (e) Calculate the time, according to observer P, that the shuttle S takes to travel from star A to star B. [2]

.....

.....

.....

.....

- (f) Identify and explain the reference frame in which the proper time for shuttle S to journey from star A to star B can be measured. [2]

.....

.....

.....


.....

(Option A continues on the following page)



(Option A continued)

5. The spacetime diagram shows the Earth frame with the worldline of a spaceship S moving away from Earth. $ct' = 0$ when $ct = 0$.



- (a) Determine the speed of the spaceship relative to Earth.

[1]

.....

.....

A flash of light sent by an Earth observer at $ct = 2.0 \text{ km}$ is directed towards the spaceship.

- (b) Estimate, using the spacetime diagram, the time in seconds when the flash of light reaches the spaceship according to the Earth observer.

[2]

.....

.....

.....

.....

(Option A continues on the following page)



(Option A, question 5 continued)

- (c) Determine the time coordinate ct' when the flash of light reaches the spaceship, according to an observer at rest in the spaceship.

[2]

.....

.....

.....

.....

.....

.....

End of Option A




32EP11

Turn over


Option B — Engineering physics

6. A student models a rotating dancer using a system that consists of a vertical cylinder, a horizontal rod and two spheres.

The cylinder rotates from rest about the central vertical axis. A rod passes through the cylinder with a sphere on each side of the cylinder. Each sphere can move along the rod. Initially the spheres are close to the cylinder.



A horizontal force of 50 N is applied perpendicular to the rod at a distance of 0.50 m from the central axis. Another horizontal force of 40 N is applied in the opposite direction at a distance of 0.20 m from the central axis. Air resistance is negligible.



(Option B continues on the following page)



- axis is approximately 20 Nm. [11]

- angular speed of 20 rad s^{-1} in a
stem. [1]

.....

.....

.....

.....

.....


- central axis. [2]

[illegible]

L

(Option B, question 6 continued)

- (d) When the system has reached its maximum angular speed, the two forces are removed. The spheres now move outward, away from the central axis.



- (i) Outline why the angular speed ω decreases when the spheres move outward. [2]

.....

.....

.....

.....

- (ii) Show that the rotational kinetic energy is $\frac{1}{2}L\omega$ where L is the angular momentum of the system. [1]

.....

.....

- (iii) When the spheres move outward, the angular speed decreases from 20 rad s^{-1} to 12 rad s^{-1} . Calculate the percentage change in rotational kinetic energy that occurs when the spheres move outward. [2]

.....

.....

.....

.....

.....

.....

(Option B continues on the following page)



(Option B, question 6 continued)

- (e) Outline one reason why this model of a dancer is unrealistic. [1]

.....


.....

(Option B continues on the following page)




(Option B continued)

7. A frictionless piston traps a fixed mass of an ideal gas. The gas undergoes three thermodynamic processes in a cycle.



The initial conditions of the gas at A are:

volume = 0.330 m^3
 pressure = 129 kPa
 temperature = 27.0°C



Process AB is an isothermal change, as shown on the pressure volume (pV) diagram, in which the gas expands to three times its initial volume.

- (a) Calculate the pressure of the gas at B.

[2]

.....

.....

.....

.....

(Option B continues on the following page)



(Option B, question 7 continued)

The gas now undergoes adiabatic compression BC until it returns to the initial volume. To complete the cycle, the gas returns to A via the isovolumetric process CA.

- (b) Sketch, on the pV diagram, the remaining two processes BC and CA that the gas undergoes. [2]
- (c) Show that the temperature of the gas at C is approximately 350°C . [2]

.....

.....

.....

.....

.....

.....

.....

- (d) Explain why the change of entropy for the gas during the process BC is equal to zero. [1]

.....

.....

- (e) Explain why the work done by the gas during the isothermal expansion AB is less than the work done on the gas during the adiabatic compression BC. [1]

.....

.....

- (f) The quantity of trapped gas is 53.2 mol. Calculate the thermal energy removed from the gas during process CA. [2]

.....

.....

.....

.....

End of Option B




32EP17

Turn over

Option C — Imaging

8. An object is placed in front of a concave mirror with the focal point f as shown.



- (a) Construct a ray diagram to locate the position of the image produced.

[2]

(Option C continues on the following page)



(Option C, question 8 continued)


(b) Describe the features of the image produced.

[1]

.....

.....

(c) Parallel light rays are incident on a spherical concave mirror as shown.



State the problem illustrated by the diagram and how it is corrected in reflecting telescopes.

[2]

.....

.....

.....


.....

(Option C continues on the following page)



(Option C continued)

9. The eye of an observer has a near point of 25 cm. A pencil is placed at the near point. A convex lens of focal length 8 cm is then placed between the pencil and the observer as shown. The pencil is positioned at the focal point of the lens.




- (a) Determine the angular magnification of the lens when the image of the pencil is viewed at infinity. [1]

.....

.....

- (b) A student increases the magnification of the pencil by using two 8 cm focal length convex lenses placed 25 cm apart. The pencil is placed 14 cm from one of the lenses.



(Option C continues on the following page)



(Option C, question 9 continued)

- (i) Show that the magnitude of the magnification of the pencil produced by the lens closest to the pencil is approximately 1.3. [2]

.....

.....

.....

.....

- (ii) Calculate the total magnification observed by the student using the two lenses as shown. [2]

.....

.....

.....

.....

- (c) The two 8 cm focal length convex lenses are now used to construct a telescope in normal adjustment. The diameter of the lenses is much greater than the diameter of the pupil of the eye. State, compared with the naked eye,

- (i) **one** advantage of using this telescope for astronomical observations. [1]

.....

.....

- (ii) **one** disadvantage of using this telescope for astronomical observations. [1]

.....

.....

(Option C continues on page 23)



Please do not write on this page.

Answers written on this page
will not be marked.



(Option C, question 9 continued)

- (d) Describe how international collaboration can improve the quality of the image of radio array telescopes.

[2]

.....

.....

.....

.....

(Option C continues on the following page)

.....

.....

.....

.....



32EP23

Turn over

(Option C continued)

10. Signals in an optic fibre require amplification when intensity levels in the fibre have fallen to 1.5% of the original signal. A light signal of initial intensity I_0 is sent down the optic fibre.

- (a) The fibre has an attenuation per unit length of 0.30 dB km^{-1} . Deduce that the length of the fibre is approximately 60 km before the signal requires amplification. [2]


.....

.....

.....

.....

A signal is sent down a 27 km step-index fibre and received according to the intensity–time graph below.



- (b) Calculate the refractive index of the fibre. [2]

.....

.....

.....

.....

(Option C continues on the following page)



(Option C, question 10 continued)

- (c) Discuss how using a graded-index fibre could reduce waveguide dispersion. [2]

.....

.....

.....

.....

End of Option C



32EP25

Turn over

Option D — Astrophysics

11. (a) The Ghost of Jupiter is a nebula.

- (i) Outline what is meant by a nebula.

[1]

.....

.....

- (ii) Astrophysicists have deduced the nature of this nebula from Earth. Outline how they can make these deductions.

[1]

.....

.....

- (b) Star X and star Y are in our own galaxy. They appear to move with respect to very distant stars when viewed from Earth during a six-month period. The following data are provided.

	Parallax angle	Apparent brightness
Star X	0.019 arc-second	$8.4 \times 10^{-9} \text{ Wm}^2$
Star Y	0.038 arc-second	$3.1 \times 10^{-9} \text{ Wm}^2$

- (i) Deduce which star will appear to move more.

[2]

.....

.....

.....

.....

- (ii) Calculate, in m, the distance to star X.

[1]

.....

.....

(Option D continues on the following page)



(Option D, question 11 continued)

- (iii) Determine the ratio $\frac{\text{luminosity of star X}}{\text{luminosity of star Y}}$.

[2]

.....

.....

.....

.....

.....

.....

.....

(Option D continues on the following page)




32EP27

Turn over

(Option D continued)

12. Three stars A, B and C are labelled on the Hertzsprung–Russell (HR) diagram. L_{\odot} is the Luminosity of the sun.



- (a) State the main element that is undergoing nuclear fusion in star C.

[1]

.....

.....

- (b) Explain why star B has a greater surface area than star A.

[2]

.....

.....

.....

.....

(Option D continues on the following page)



(Option D, question 12 continued)

- (c) White dwarfs with similar volumes to each other are shown on the HR diagram.

Construct a line, on the HR diagram, to show the possible positions of other white dwarf stars with similar volumes to those marked on the HR diagram.

[2]

- (d) Some stars on the HR diagram are likely to evolve into neutron stars.

Outline why the radius of a neutron star reaches a stable value.

[2]

.....

.....

.....

.....

(Option D continues on the following page)



32EP29

Turn over

(Option D continued)

13. Galaxy D has a redshift $z = 0.13$.

- (a) Calculate, in Mpc, the distance to D using a Hubble constant value of $73 \text{ km s}^{-1} \text{ Mpc}^{-1}$. [2]

.....

.....

.....

.....

- (b) A Hubble constant value of $73 \text{ km s}^{-1} \text{ Mpc}^{-1}$ gives an age of the universe to be 13.4×10^9 years when assuming a constant rate of expansion has occurred.

- (i) Determine in years, the age of the universe when the light detected on Earth now was originally emitted from D. [3]

.....

.....

.....

.....

.....

.....

- (ii) Evidence based on observations of type Ia supernovae affects the result in (b)(i). State the relevant conclusion from these observations. [1]

.....

.....

End of Option D

